

Lacustrine otoliths fauna from the Miocene sediments of Vračević (Western Serbia)

Katarina Bradić-Milinović, Ljupko Rundić, Werner Schwarzhans



Дигитални репозиторијум Рударско-геолошког факултета Универзитета у Београду

[ДР РГФ]

Lacustrine otoliths fauna from the Miocene sediments of Vračević (Western Serbia) | Katarina Bradić-Milinović, Ljupko Rundić, Werner Schwarzhans | 9th International workshop on Neogene of Central and South-East Europe, 6th – 9th June, Tuzla, Bosnia and Herzegovina | 2022 | |

10.51558/2303-5161..2022.1.1.21

<http://dr.rgf.bg.ac.rs/s/repo/item/0007052>

Дигитални репозиторијум Рударско-геолошког факултета Универзитета у Београду омогућава приступ издањима Факултета и радовима запослених доступним у слободном приступу. - Претрага репозиторијума доступна је на www.dr.rgf.bg.ac.rs

The Digital repository of The University of Belgrade Faculty of Mining and Geology archives faculty publications available in open access, as well as the employees' publications. - The Repository is available at: www.dr.rgf.bg.ac.rs



LACUSTRINE OTOLITHS FAUNA FROM THE MIOCENE SEDIMENTS OF VRAČEVIĆ (WESTERN SERBIA)

Katarina Bradić-Milinović¹, Ljupko Rundić², Werner Schwarzhans³

Here we present a rich assemblage of fish otoliths from the late Badenian to early Sarmatian (MN7+8 zone), which were collected along the Grabovac stream in the village of Vračević. They were obtained from freshwater thin-bedded marly deposits of the Vračević part of the Serbian Lake System in the Valjevo-Mionica Basin in western Serbia. This basin covers an area of 350 km² and represents the western part of the so-called Valjevo-Mionica-Belanovica Graben that had formed during the Ottnangian-Karpatian, and later became inverted (Marović et al, 2007).

The stratigraphic position of the Vračević locality is controversial. Based on terrestrial gastropods Neubauer et al. (2016) considered, these freshwater sediments as Sarmatian s.s. in age. According to terrestrial micromammals the sites represent the MN7+8 zone (Marković, 2003). Recently, a shallow well drilled at the hill close to the Grabovac stream, gave evidence of the presence of the marine to brackish Sarmatian s.s. and brackish Pannonian sediments (Bradić-Milinović et al., 2018).

The otoliths correlate well with the fauna from the early to middle Miocene of Klinci (Bradić-Milinović et al., 2019). The fish fauna from Klinci represents an endemic paleobioprovince characteristic to the ancient Serbian Lake System (SLS). We observed a similar type of fish fauna in both localities, which indicates that they have connected through time during the existence of the Valjevo-Mionica Basin. The freshwater community of goby fishes from the early to early middle Miocene of southeastern Europe documents the presence of a lost Miocene freshwater goby fish fauna, before the unrelated Ponto-Caspian fish fauna of today expanded into the freshwater environments of its borderland (Bradić-Milinović et al., 2019).

We identified 9 different species, of which two are new and three remain in open nomenclature: *Aphanius jeani*, *Aphanolebias* n.sp., *Klincigobius andjelkovicae*, *Klincigobius* n.sp., *Klincigobius serbiensis*, *Klincigobius* sp., *Toxopyge campylus*, *Toxopyge* sp. and *Ponticola* sp. (Bradić-Milinović et al., submitted).

Based on the composition of the fish fauna in the Vračević sediments, we concluded that it could be older in age than Sarmatian s.s., but younger than the fauna from Klinci (Bradić-Milinović et al., submitted). In fact it so far represents the last occurrence of this lost freshwater goby fauna.

¹ Geological Survey of Serbia, Rovinjska 12, 11 000 Belgrade, Serbia.
Corresponding author: katarina.bradicmilinovic@gzs.gov.rs

² University of Belgrade, Faculty of Mining and Geology, Department of Regional Geology, Kamenička 6, 11 000 Belgrade, Serbia; e-mail: ljupko.rundic@rgf.bg.ac.rs

³ Natural History Museum of Denmark, Zoological Museum, Universitetsparken 15, 2100 København, Denmark; e-mail: wwschwarz@aol.com.




Figure 1- The lost early to middle Miocene freshwater goby association of central and southeastern Europe and the role of the Klinci and Vračević faunas in it.

REFERENCES

- Bradić–Milinović, K., Rundić, Lj. & Bojić, Z. (2018). A contribution for stratigraphy of the Miocene of Vračević (Valjevo–Mionica Basin). Proceedings of XVII Congress of the geologists of Serbia, 1, 110–114.
- Bradić–Milinović, K., Ahnelt, H., Rundić, Lj. & Schwarzhans, W. (2019). The lost fresh water goby fish fauna (Teleostei, Gobiidae) from the early Miocene of Klinci (Serbia). Swiss Journal of Palaeontology, 138, 285–315.
- Bradić–Milinović, K., Rundić, Lj. & Schwarzhans W. (submitted). Middle Miocene freshwater otoliths from the Vračević lake (Serbian Lake System). Annales Géologiques de la Péninsule Balkanique
- Marović M., Toljić M., Rundić Lj. & Milivojević J. (2007). Nealpine tectonics of Serbia. Serbian Geological Society, 1–82.
- Marković, Z. (2003). The Miocene small mammals of Serbia, a review. In: Reumer, J. W. F. & Wessels, W. (ed.) – Distribution and migration of Tertiary mammals in Eurasia. A volume in honour of Hans de Bruijn – DEINSEA 10, 393–398.
- Neubauer, T., A., Harzhauser, M., Mandić, O. & Jovanović, G. (2016). The late middle Miocene non–marine mollusk fauna of Vračević (Serbia): filling a gap in Miocene land snail biogeography. Bulletin of Geosciences, 91(4), 731–778.

Acknowledgment: This study was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract on realization and financing of Scientific Research Work NIO in 2022 No.451-03-68/2022-14/200126)