

Arsenic in tape waters of the south Pannonian basin (Serbia) and arsenic risk assessment

Petar Papić, Marina Ćuk, Maja Todorović, Jana Stojković, Bojan Hajdin, Nebojša Atanacković, Dušan Polomčić



Дигитални репозиторијум Рударско-геолошког факултета Универзитета у Београду

[ДР РГФ]

Arsenic in tape waters of the south Pannonian basin (Serbia) and arsenic risk assessment | Petar Papić, Marina Ćuk, Maja Todorović, Jana Stojković, Bojan Hajdin, Nebojša Atanacković, Dušan Polomčić | Polish Journal of Environmental Studies | 2012 | 21 | 6

<http://dr.rgf.bg.ac.rs/s/repo/item/0002196>

Дигитални репозиторијум Рударско-геолошког факултета Универзитета у Београду омогућава приступ издањима Факултета и радовима запослених доступним у слободном приступу. - Претрага репозиторијума доступна је на www.dr.rgf.bg.ac.rs

The Digital repository of The University of Belgrade Faculty of Mining and Geology archives faculty publications available in open access, as well as the employees' publications. - The Repository is available at: www.dr.rgf.bg.ac.rs

certain samples of drinking water collected in several towns within the riparian zone of the Danube measured elevated arsenic concentrations. In South Banat the ammonium ion concentrations are 0.72-2.75 mg/L, while the MPC for this ion in drinking water is 0.5 mg/L [24].

Arsenic Occurrence in Groundwater of Bačka District

Bačka generally obtains its drinking water supply from centralized public water supply systems. Rural settlements rely on local groundwater sources. With regard to the cation




Fig. 3. Histogram of arsenic in groundwater used for drinking water supply in Banat.




Fig. 4. Histogram of arsenic concentrations in groundwater used for drinking water supply in Bačka.




Fig. 5. Arsenic concentrations in Srem groundwaters.

composition, sodium dominates all locations and all water-bearing media, except the Quaternary sediments in the extended area of Subotica, where the cation composition was primarily Ca-Mg [21]. The average arsenic concentration in Bačka groundwaters was 48 µg/L (Fig. 4).

The largest city in this district is Subotica (population 140,000), and this is the only city with a water treatment plant. The extended area obtains its drinking water supply from 13 groundwater sources. Based on the analyses, the average concentration of arsenic at these groundwater sources was 110 µg/L. The presence of the ammonium ion and absence of nitrates in groundwater of the test area suggests a reducing medium that creates favorable conditions for arsenic migration in groundwater. Due to the low solubility of iron oxides, hydroxides, and carbonates in groundwater, their concentrations in groundwater were found to be rather low. The iron concentration in the groundwater used for drinking water supply in Subotica measured from 0.04 to 0.96 mg/L.

Arsenic Occurrence in Groundwater of Srem District

The Srem District of Vojvodina features groundwater of good quality, with arsenic concentrations falling within the allowed range and 77% of samples having arsenic concentrations below 10 µg/L (Fig. 5). The most frequent causes of physicochemical non-compliance in Srem are concentrations of iron and manganese higher than MPC [19].

In the Srem District, the mineral content of groundwater was found to be lower than in other parts of Vojvodina. Total hardness measured from 5°dH to 17°dH. Concentrations of organic substances (KMnO₄ demand) were below 12 mg/L, while Fe concentrations measured about 0.3 mg/L. The public water supply is generally provided by tapping aquifers up

to a depth of 250 m, which meet drinking water standards. Depths below 250 m exhibited elevated total mineralization [19].

Arsenic Risk Assessment by the Drinking Water Pathway

The end result of the assessment of groundwaters used for drinking water supply in the South Pannonian Basin should point to the importance of finding an alternative water supply for most exposed populations. This paper gives a prognostic hydrochemical map showing arsenic concentrations (Fig. 6) and assessment of the risks incurred through drinking water. According to the Provincial Secretariat of Science and Technology Development, more than 600,000 inhabitants of Banat and Bačka (or some 40% of the population of Vojvodina) obtain drinking water supply with arsenic concentrations exceeding maximum allowable concentrations in drinking water (10 µg/L) [19]. However, the actual number of the people in Vojvodina exposed to As may be lower due to the use of bottled drinking water and/or the use of private or community shallow wells, which are assumed to be As free. On the other hand, the residents of some of the villages in the study area may be exposed to drinking water with As concentrations that could lead to cancer. In comparison with studied in West Bengal, India, more than 6 million people are exposed to arsenic through their drinking water, among which approximately 300,000 people manifest signs of chronic arsenicosis [36]. It was found (in West Bengal) that it produces various systemic manifestations such as chronic lung disease, characterized by chronic bronchitis, chronic obstructive and/or restrictive pulmonary disease, and bronchiectasis;

liver diseases, such as non-cirrhotic portal fibrosis; polyneuropathy; peripheral vascular disease; hypertension; nonpitting edema of feet/hands; conjunctival congestion; weakness; and anemia [37]. In Serbia the impact of arsenic on human health is insufficiently researched. The Institute of Public Health of Serbia sees association between exposure to arsenic in drinking water and the occurrence of type 2 diabetes in the Middle Banat region. The study has shown a significantly higher odds ratio and higher standardized incidence rates for the occurrence of type 2 diabetes in the population exposed to arsenic, compared to the unexposed population in Serbia [38].

The highest safe standard for cancer risk is 1 per 10,000 and the lowest standard 1 per 1,000,000 [39]. Furthermore, in the study area the cancer risk indices (R) ranged from 0.13 to 2.6 per 1,000 persons. To assess the health effects of chronic As exposure it would be necessary to accurately characterize residents' consumption patterns, namely to track changes in the sources of drinking water, duration of use, and drinking rate.

Central Banat and North Bačka featured the highest arsenic concentrations in groundwaters used for drinking water supply. At most water supply systems in South Banat arsenic concentrations remained below MPC, while levels in excess of 10 µg/L were recorded in populated areas in the riparian zone of the Danube. The Srem District generally exhibited the best quality of groundwater used for drinking water supply, with arsenic concentrations remaining below 10 µg/L.

The natural baseline concentrations of As in the groundwater defined during the hydrochemical study and presented in this paper were used for As risk assessment. The results obtained indicated that residents who consume raw




Fig. 6. Prognostic hydrochemical map with arsenic zoning of cities in Vojvodina.

